arXiv:1203.1275v1 [math.PR] 6 Mar 2012

Moments and Classification for Conjugation-Invariant Rotations and Fake Uniformity in the Stochastic Radon Transform

Christian Rau*
Adjunct Research Fellow
School of Mathematical Sciences, Monash University
Clayton VIC 3800, Australia

05 March 2012

Abstract

We consider a generalisation of the stochastic Radon transform, introduced for an inverse problem in tomography by Panaretos. Specifically, we allow the distribution of the three-dimensional rotation in the statistical model of that work to be different from Haar measure, and to possess only the weaker symmetry property of conjugation-invariance. As a preparation we study, based on simple geometric ideas, the relationship between moments of two univariate densities describing conjugation-invariant rotations. For the class of rotations introduced by León, Massé and Rivest, the generalised stochastic Radon transform turns out to have a 'fake uniformity' property. We also derive the binary Bayes classifier for rotations whose distributions are deterministic shifts of a common conjugation-invariant distribution.

Keywords: Classification, fake uniformity, inverse problems, moments, Radon transform, rotations, shape theory.
AMS 2000 Subject Classification: Primary 60D05; Secondary 62H05, 62H30.

1 Introduction

Random rotations in Euclidean space $\mathbb{R}^d$ ($d \geq 2$) arise in a variety of contexts in stochastic geometry, especially in the construction of other stochastic models, for example random hyperplane processes; see the monograph [13]. In these contexts, the rotation distribution is usually taken as the Haar probability measure $\mu = \mu_d$ on the d-dimensional rotation group $SO(d)$. Recall that μ

---

[*] Author's email address: christianrau88@gmail.com

is characterised by the property that $\mu(QA) = \mu(AQ) = \mu(A)$ for any $Q \in SO(d)$ and any measurable $A \subseteq SO(d)$, where $SO(d)$ is endowed with its Borel σ-field, $QA = \{QP : P \in SO(d)\}$, and AQ is defined analogously. See [13, Ch. 13] for more information and one of several constructions of Haar measure on $SO(d)$. Most applications concern the case $d = 3$, not only in stochastic geometry but also statistics. Two important instances are a statistical model for random orientations of limbs of experimental subjects performing drilling tasks, studied in [4], and a model for tomographic imaging introduced in [7] and further studied in [8]. In [4], a class of rotation distributions in $\mathbb{R}^d$ was introduced that was shown to be amenable to exact computations of several key quantities including first and second moments. In [7], the goal was to estimate the shape (as defined in that paper) of a particle imaged with cryo-electron microscopy, where the orientation of the particle is not only random but also unknown, its orientation being governed by μ. As was shown in that paper, this last assumption is necessary in order to produce a well-defined estimator of the shape of the particle.

The basis of the present paper is a synthesis of [4] and [7]. Our focus is on probabilistic rather than statistical aspects of the model. Specifically, we study a probability model that arises when the rotation distribution in the stochastic Radon transform introduced in [7] is not Haar measure, but only possesses the more general symmetry property of conjugation-invariance, whose definition we recall in Section 2. Our results will be more general than what we need for the tomography application, and in particular, we will obtain several new results relating to three-dimensional rotation distributions. We also derive the Bayes classification rule with two classes which are shifts of a common conjugation-invariant distribution, and examine the behaviour for small shifts. For the case of two multivariate Normal distributions on $\mathbb{R}^d$ with equal covariance matrices, a corresponding result is well known; see [10, pp. 21–22]. As far as we are aware, our approach is essentially new, although there is some similarity in some instances with the geometric reasoning in [3], which studied the distribution of the projected shape under Haar measure.

Our two main results, namely Theorems 1 and 2 in Sections 4 and 5 respectively, yield two different ways of quantifying the effect of moving away from the Haar measure situation. In relation to the first of these results, a finding of particular interest is what we call a *fake uniformity* property, ‘uniform’ being taken here and below as a synonym for Haar measure μ (see the discussion around Definition 1 in Section 4). The second result quantifies how well one may distinguish two distributions for a shift of ‘modal’ angle, as explained later; as expected, we find better accuracy for stronger deviation of the common ‘centred’ distribution from μ (see the discussion around (12) in Section 5). We shall not attempt to develop a variational approach using distances between rotation distributions as was done in [9], as this seems a secondary matter to the phenomena we study. Our results also provide a new interpretation of the concentration parameter in statistical models for random rotations. Finally, our results may provide a foundation for statistical tests that are based on the quantities we consider. It seems clear that the fake uniformity property is a problem for the most basic of such tests, namely that for uniformity of the rotation.

Section 2 contains background on rotations, as well as notation. Section 3 investigates the connection between two univariate distributions which describe conjugation-invariant rotations. Section 4 contains the main result on the stochastic Radon transform for what we call shift-symmetric rotations, along with an example that reveals the fake uniformity property and motivates us to define it separately. Finally, in Section 5 the Bayes classifier is derived.

2 Notation and Background

All random variables and random elements on other state spaces are taken to reside on a common probability space $(\Omega, \mathscr{F}, \mathrm{P})$, with $\mathrm{E}(\cdot)$ denoting the corresponding expectation operator.

We shall be concerned not only with random variables on $\mathbb{R}$, but also on the sphere in three dimensions $\mathbb{S}^2 := \{x \in \mathbb{R}^3 : \|x\| = 1\}$, and on $SO(3)$. If $\boldsymbol{T}$ denotes a random element in either of these three spaces, we shall denote its density by $f_{\boldsymbol{T}}$, it being understood that the dominating measure for this density is ordinary Lebesgue measure, normalised Lebesgue surface measure σ on $\mathbb{S}^2$, or Haar probability measure μ on $SO(3)$ respectively. As done above, we use boldfacing to distinguish deterministic and random rotations, but will use uppercase and lowercase letters for vectors and scalars as is customary. All vectors are regarded as column vectors. We write $v_i = v \cdot e^{(i)}$ for $v \in \mathbb{R}^3$, where the dot denotes the usual inner product and $e^{(i)}$ is the ith column of the 3×3 identity matrix, for $i = 1, 2, 3$. The matrix $\mathrm{diag}(a_1, a_2, a_3)$ is the 3×3 diagonal matrix with entries a_1, a_2, a_3 along its main diagonal. The Beta integral is

$$\mathrm{B}(p, q) = \int_0^1 x^{p-1}(1-x)^{q-1}\,, \qquad p, q > 0\,,$$

and $\mathrm{Beta}(p, q)$ is the Beta distribution with parameters $p, q > 0$. Recall that if $W \sim \mathrm{Beta}(p, q)$ then

$$\mathrm{E}(W^r) = \frac{(p)_r}{(p+q)_r} \qquad \text{for any integer } r \geq 0, \tag{1}$$

where $(x)_n = \Gamma(x+1)/\Gamma(x-n+1)$ is the Pochhammer symbol and $\Gamma(\cdot)$ is the Gamma function. We use the notation $\mathrm{E}(X; B) := \mathrm{E}(X 1_B)$ for a scalar random variable X and a measurable set $B \subset \mathbb{R}$, where 1_B is the indicator function of B.

Recall that any rotation $P \in SO(3)$ (except if it belongs to the null set comprising the identity and rotations by an angle π) may be expressed uniquely in terms of the rotation axis $u \in \mathbb{S}^2$

and rotation angle $\theta \in [0, \pi)$, as follows (see [11, pp. 188–189] or [12, p. 99]):

$$P = \cos(\theta) I + \sin(\theta) \mathcal{S}(u) + (1 - \cos(\theta)) u u^T \,, \quad \text{where} \quad \mathcal{S}(a) = \begin{pmatrix} 0 & -a_3 & a_2 \\ a_3 & 0 & -a_1 \\ -a_2 & a_1 & 0 \end{pmatrix} . \tag{2}$$

We assume throughout that the letter $\boldsymbol{R}$ denotes a random three-dimensional rotation which is *conjugation-invariant*, meaning that is satisfies $\boldsymbol{R} \overset{\mathrm{d}}{=} Q^T \boldsymbol{R} Q$, where $\overset{\mathrm{d}}{=}$ denotes equality in distribution; then automatically $\boldsymbol{R}^T \overset{\mathrm{d}}{=} \boldsymbol{R}$. The terminology here follows [12]. We say that $\boldsymbol{R}$ (or its distribution) belongs to $\mathcal{I}_s$. This definition for $\mathcal{I}_s$ is somewhat less restrictive than the one given in [11, p. 190]. In the context of random $\boldsymbol{P}$ at (2), one has $f_{\boldsymbol{P}} \in \mathcal{I}_s$ if and only if $U \sim \sigma_2$ (cf. [11, p. 191]). In the sequel, when referring to random elements U and Θ, we always assume them to be defined through $\boldsymbol{P}$ via the bijection (up to a null set) $\boldsymbol{P} \leftrightarrow (U, \Theta)$.

To date, only few rotation distributions have been studied in the literature. We shall consider the distribution introduced in [4], referred to below as the Cayley-LMR distribution:

$$f_{\mathrm{CAY}(M,\kappa)}(P) = \frac{\sqrt{\pi}\,\Gamma(\kappa+2)}{2^{2\kappa}\Gamma(\kappa+1/2)} (1 + \operatorname{tr}(P M^T))^{\kappa} \,, \tag{3}$$

and the symmetric matrix Fisher-von Mises distribution, whose density is proportional to $\exp\{\kappa(PM^T)\}$; see [5, pp. 289ff] for more information on the latter distribution. In both cases, $\kappa \geq 0$ is a dispersion parameter, with $\kappa = 0$ corresponding to the case $f_{\boldsymbol{P}} = f_{\boldsymbol{R}} = \mu$, and M may be regarded as a modal matrix. We shall make the minimal assumption that our rotation distributions are parametrised by modal matrices in the sense of a group model, see [9, p. 653], and a dispersion parameter κ, with the case $\kappa = 0$ corresponding to Haar measure μ. They are allowed to have other parameters as well.

We shall on several occasions consider the random variable

$$X := \frac{1 + \cos\Theta}{2} \qquad \left(= \cos^2\left(\frac{\Theta}{2}\right) \right) \tag{4}$$

and its (central) moments $\{\rho_r = \mathrm{E}(X^r),\, r \geq 0\}$. For the Cayley-LMR distribution, $X \sim \mathrm{Beta}(\kappa + 1/2, 3/2)$ ([4], p. 425) and thus by (1),

$$\rho_1(\kappa) = \frac{\kappa + 1/2}{\kappa + 2} \,, \qquad \rho_2(\kappa) = \frac{(\kappa+1/2)(\kappa+3/2)}{(\kappa+2)(\kappa+3)} \,. \tag{5}$$

It will become clear later why we regard these as functions of κ.

Finally, we introduce some convenient terminology. For a random rotation $\boldsymbol{P}$ which is such that there is a conjugation-invariant rotation $\boldsymbol{R} \in SO(3)$ and a deterministic rotation $M \in SO(3)$ such that $\boldsymbol{P} M^T \overset{\mathrm{d}}{=} \boldsymbol{R}$, we write $\boldsymbol{P} \in \mathcal{I}_s^M$. We call $\boldsymbol{Q}$ *shift-symmetric*, and write $\boldsymbol{Q} \in \mathcal{I}_s^+$, if there exists an M such that $\boldsymbol{Q} \in \mathcal{I}_s^M$. The same terminology applies to the underlying distributions of the random variables.

3 Relation between two distributions and moments

In this section we investigate the connection between two key distributions derived from a three-dimensional rotation. The first proposition shows that if $f_{\boldsymbol{R}} \in \mathcal{I}_s$ then the density $f_{\boldsymbol{R}e^{(3)}}$ is simply that of

$$Z = \left(e^{(3)}\right)^T \boldsymbol{R} e^{(3)} \,. \tag{6}$$

In the proof and in several later instances we shall need the following simple result on integration on the sphere, which is a special case of [1, Proposition 9.1.2, p. 189]:

Lemma 1. Let f be an integrable function on $\mathbb{S}^2$ which is zonal, that is, $f(x) = F(x_3)$, where F is a function defined on $[-1, 1]$. Then

$$\int_{\mathbb{S}^2} f(x)\, \sigma(\mathrm{d}x) = \frac{1}{2} \int_{-1}^{1} F(t)\, \mathrm{d}t \,.$$

In particular, if $U \sim \sigma_2$ then $U_3 \sim$ Uniform$[-1, 1]$.

Proposition 1. $f_{\boldsymbol{R}e^{(3)}}(w) = f_Z(w_3)$. Moreover, $f_{\boldsymbol{R}e^{(3)}}$ determines $f_{\boldsymbol{R}}$ uniquely.

Proof. Consider realisations $R = \boldsymbol{R}(\omega_0)$, $z = Z(\omega_0)$, $u = U(\omega_0)$, and $\theta = \Theta(\omega_0)$. By the law of cosines for spherical triangles (see [2, p. 54]), we have

$$z = \cos\left(\angle(e^{(3)}, Re^{(3)})\right) = \cos^2\left(\angle(u, Re^{(3)})\right) + \sin^2\left(\angle(u, Re^{(3)})\right)\cos\theta = u_3^2 + (1 - u_3^2)\cos\theta \,, \tag{7}$$

where $\angle(v, w) = \arccos(v \cdot w) \in [0, \pi)$ is the angle between the vectors $v, w \in \mathbb{S}^2$. Hence $\cos\Theta = (Z - U_3^2)/(1 - U_3^2)$ almost surely. The equality of the densities follows from Lemma 1, cf. also the remark in [1, p. 189]. Now (2) shows that the distribution of $\boldsymbol{R}$, and thus also that of $\boldsymbol{R}e^{(3)}$, is determined uniquely. □

Corollary 1. If $\boldsymbol{Q} \in \mathcal{I}_s^M$ with M such that $Me^{(3)} = e^{(3)}$, then $f_{\boldsymbol{Q}e_3}(w_3) = f_Z(w_3)$.

Proof. Follows from the fact that [9, p. 653] $f_{\boldsymbol{Q}e_3} = f_{\boldsymbol{R}e_3} \circ \Pi^{-1}$ for any $\Pi \in SO(3)$ with $Me^{(3)} = \Pi e^{(3)}$. □

Proposition 2.

$$f_Z(s) = \frac{1}{2\sqrt{2}} \int_0^{\frac{1+s}{2}} \frac{f_X(x)}{\sqrt{(1 + s - 2x)(1 - x)}}\, \mathrm{d}x \,, \qquad -1 \le s \le 1 \,.$$

Proof. By Lemma 1, (7) and Fubini's theorem,

$$\begin{aligned}
\mathrm{P}(Z \le s) &= \frac{1}{2}\int_{-1}^{1} \mathrm{P}(Z \le s \mid U_3 = t)\,\mathrm{d}t = \frac{1}{2}\int_{-1}^{1} \mathrm{P}\big\{t^2 + (1-t^2)\cos\Theta \le s\big\}\,\mathrm{d}t \\
&= \frac{1}{2}\int_{-1}^{1} \mathrm{P}\left\{X \le \frac{1+s-2t^2}{2(1-t^2)}\right\}\mathrm{d}t = \frac{1}{2}\int_{-1}^{1}\int_{0}^{\frac{1+s-2t^2}{2(1-t^2)}} f_X(x)\ \mathrm{d}x\,\mathrm{d}t \\
&= \frac{1}{\sqrt{2}}\int_{0}^{\frac{1+s}{2}} \sqrt{\frac{1+s-2x}{1-x}}\, f_X(x)\,\mathrm{d}x\,,
\end{aligned}$$

and differentiation with respect to s completes the proof. □

Remark 1. Conversely, it is readily shown that

$$f_X(s) = \frac{1}{2}\int_{-1}^{1} f_Z(2s - 2st + 2t^2 - 1)\,\mathrm{d}t\,, \qquad 0 \le s \le 1\,.$$

The relationship between the densities in Proposition 2 will now be used to relate the corresponding moments up to some finite order. Consider the vector $\tau = (\tau_j)_{j=0}^{m}$, $m \ge 1$, where $\tau_j = \mathrm{E}(Z^j)$. Lemma 1 and Proposition 2 yield for $k \ge 1$, by partial integration,

$$\begin{aligned}
\tau_k &= \frac{1}{2\sqrt{2}}\int_{-1}^{1} t^k\,\mathrm{d}t \int_{0}^{\frac{1+t}{2}} \frac{f_X(x)}{\sqrt{(1+t-2x)(1-x)}}\,\mathrm{d}x \\
&= \frac{1}{2\sqrt{2}}\int_{0}^{1} \frac{f_X(x)}{\sqrt{1-x}}\,\mathrm{d}x \int_{2x-1}^{1} \frac{t^k}{\sqrt{1+t-2x}}\,\mathrm{d}t \\
&= 1 - \frac{k}{\sqrt{2}}\int_{0}^{1} \frac{f_X(x)}{\sqrt{1-x}}\,\mathrm{d}x \int_{2x-1}^{1} t^{k-1}\sqrt{1+t-2x}\,\mathrm{d}t\,. \qquad (8)
\end{aligned}$$

Another partial integration gives, for any $k \ge 1$ and $-1 \le x \le 1$,

$$\begin{aligned}
G_k(x) &:= \int_{2x-1}^{1} t^{k-1}\sqrt{1+t-2x}\,\mathrm{d}t = \frac{4\sqrt{2}}{3}(1-x)^{3/2} - \frac{2(k-1)}{3}\int_{2x-1}^{1} t^{k-2}(1+t-2x)^{3/2}\,\mathrm{d}t \\
&= a_k(x) + b_k(x)\,G_{k-1}(x)\,,
\end{aligned}$$

where $G_0 \equiv 0$ and

$$a_k(x) = \frac{4\sqrt{2}}{2k+1}\,(1-x)^{3/2}\,, \qquad b_k(x) = \frac{2(k-1)}{2k+1}\,(2x-1)\,.$$

By induction,

$$G_k(x) = \sum_{t=1}^{k}\left(\prod_{s=k-t+2}^{k} b_s(x)\right) a_{k-t+1}(x)\,, \qquad k \ge 1\,,$$

from which it follows that

$$\begin{aligned}
G_k^0(x) := \frac{G_k(x)}{\sqrt{1-x}} &= 4\sqrt{2}\,(1-x)\sum_{t=1}^{k}\left[\{2(2x-1)\}^{t-1}\frac{(k-1)!}{(k-t)!}\prod_{s=k-t+1}^{k}\frac{1}{2s+1}\right]\\
&= 2\sqrt{2}\,\frac{(k-1)!}{\Gamma(k+3/2)}\,(1-x)\sum_{t=1}^{k}\frac{\Gamma(k-t+3/2)}{(k-t)!}\,(2x-1)^{t-1}\\
&= 2\sqrt{2}\,\frac{(k-1)!}{\Gamma(k+3/2)}\,(1-x)\sum_{s=0}^{k-1}\frac{(2x)^s}{s!}\sum_{t=s+1}^{k}(-1)^{t-1-s}\frac{(t-1)!}{(t-1-s)!}\,\frac{\Gamma(k-t+3/2)}{(k-t)!}\\
&= 2\sqrt{2}\,\frac{(k-1)!}{\Gamma(k+3/2)}\,(1-x)\sum_{s=0}^{k-1}\frac{(2x)^s}{s!}\sum_{t=0}^{k-s-1}(-1)^t\,\frac{(t+s)!}{t!}\,\frac{\Gamma(k-t-s+1/2)}{(k-t-s-1)!}\,.
\end{aligned}$$

It seems impossible to reduce this further. We have

$$G_1^0(x) = \frac{4\sqrt{2}}{3}(1-x)\,, \qquad G_2^0(x) = \frac{4\sqrt{2}}{15} + \frac{4\sqrt{2}}{5}\,x - \frac{16\sqrt{2}}{15}\,x^2\,,$$

which combined with (8) gives

$$\tau_1 = -\frac{1}{3} + \frac{4}{3}\,\rho_1\,, \qquad \tau_2 = \frac{7}{15} - \frac{8}{5}\,\rho_1 + \frac{32}{15}\,\rho_2\,. \tag{9}$$

The coefficient of ρ_k in the expression for τ_k is readily seen to be equal to $k!2^{k-1}\sqrt{\pi}/\Gamma(k+3/2) \to 2$ as $k \to \infty$, by Stirling's formula.

4 Gram inversion for the stochastic Radon transform and fake uniformity

In this section we consider the stochastic Radon transform, originally introduced in [7] in the case where the rotation distribution is Haar measure μ, for conjugation-invariant rotation distributions, and will utilise results from the previous section. We first briefly repeat the main points of the stochastic Radon transform that are relevant for our own purposes; more information can be found in [7] and the references cited therein. The definition of the stochastic Radon transform in [7] is based on the projection operator Π of an essentially bounded integrable function ρ on $\mathbb{R}^3$, taken to have total mass unity and integral mean zero, with support contained in a ball of radius π centred at the origin:

$$(\Pi\{\rho\}(A))(x_1,x_2) := \int_{-\infty}^{\infty} A\rho(x_1,x_2,x_3)\,\mathrm{d}x_3\,, \qquad A \in SO(3)\,,$$

where the rotation group acts on functions on $\mathbb{R}^3$ via $(A\rho)(w) = \rho(A^{-1}w)$. For a random rotation, the map Π gives rise to a distribution P_ρ on an L^2 space of planar functions

$$P_\rho(B) = \mu\{A \in SO(3) : \Pi\{\rho\}(A) \in B\}, \tag{10}$$

where B ranges over the Borel sets in L^2 space. Note that here we consider what we may call the 'population version' of the stochastic Radon transform.

In [7] it was shown that $P_{Q\rho} = P_\rho$ for all $Q \in SO(3)$, which implies that only the *shape* of ρ, that is the orbit of ρ under the above group action, may be recovered in the estimation from sample data. This motivates the consideration of the Gram matrix. The Gram matrix of $V \in \mathbb{R}^{d\times k}$ is $\mathrm{Gram}(V) = V^T V$, where $k \geq 1$, and $d = 3$ for the original shape and $d = 2$ for its projection. It comprises a set of k 'landmarks' as its columns, and is appropriate for describing shape because $\mathrm{Gram}(AV) = \mathrm{Gram}(V)$ for any $A \in SO(3)$ or $SO(2)$ respectively.

We now present an extension of the Gram inversion formula from [7, Theorem 4.1, p. 3285] to the case of conjugation-invariant, and indeed only shift-symmetric, rotations as defined in Section 2.

Theorem 1. Let V be a $3 \times k$ matrix, $k \geq 1$. Let $H = \mathrm{diag}(1, 1, 0)$. Then for $f_{\boldsymbol{P}} \in \mathcal{I}_s^M$,

$$\int_{SO(3)} \mathrm{Gram}(HAV) f_{\boldsymbol{P}}(A)\mu(\mathrm{d}A) = V^T M^T (I - D^2) MV = \mathrm{Gram}(V) - \mathrm{Gram}(DMV),$$

where $D^2 = \mathrm{diag}\left((1-\tau_2)/2, (1-\tau_2)/2, \tau_2\right)$.

Proof. As in [7, pp. 3285–3286], one obtains from Proposition 1 and the fact that $\boldsymbol{R}^T \overset{\mathrm{d}}{=} \boldsymbol{R}$ that

$$\begin{aligned}\int_{SO(3)} \mathrm{Gram}(HAV) f_{\boldsymbol{P}}(A)\mu(\mathrm{d}A) &= V^T M^T \cdot \left[I - \mathrm{E}\left\{(\boldsymbol{R}e^{(3)})(\boldsymbol{R}e^{(3)})^T\right\}\right] \cdot MV \\ &= \mathrm{Gram}(V) - \mathrm{Gram}(DMV), \\ D^2 = \mathrm{E}\left\{(\boldsymbol{R}e^{(3)})(\boldsymbol{R}e^{(3)})^T\right\} &= \left(\int_{\mathbb{S}^2} w_i w_j f^0_{\boldsymbol{R}e^{(3)}}(w)\, \sigma_2(\mathrm{d}w)\right)_{i,j}.\end{aligned}$$

From Proposition 1 it follows that D^2 is diagonal with $\mathrm{E}\{\mathrm{tr}(D^2)\} = \mathrm{E}\{(\boldsymbol{R}e^{(3)})^T(\boldsymbol{R}e^{(3)})\} = 1$. This completes the proof. □

Remark 2. The first equation of the above proof is easily adaptable to rotations in $\mathbb{R}^d$, $d \geq 2$.

The following example reveals important consequences of the results developed so far.

Example 1. For the Cayley-LMR distribution defined at (3), from Propositions 1 and 2 we rederive for the case of rotations in $\mathbb{R}^3$ the result stated for general finite dimension in [4, Proposition 3.3, p. 419]:

$$f^{\mathrm{CAY}}_{\boldsymbol{R}e^{(3)}}(w) = 2^{-\kappa}\,(\kappa+1)\,(1+w_3)^{\kappa}\,, \qquad w \in \mathbb{S}^2\,.$$

Theorem 1 and equations (9), (5) yield

$$\int_{SO(3)} \mathrm{Gram}\{HAV\} f_{\mathrm{CAY}(M,\kappa)}(A)\mu(\mathrm{d}A) = \mathrm{Gram}(V) - \mathrm{Gram}(D^{\mathrm{CAY}}_{\kappa} MV)\,,$$
$$D^{\mathrm{CAY}}_{\kappa} = \mathrm{diag}\left(\sqrt{\frac{2(\kappa+1)}{6+5\kappa+\kappa^2}}, \sqrt{\frac{2(\kappa+1)}{6+5\kappa+\kappa^2}}, \sqrt{\frac{2+\kappa+\kappa^2}{6+5\kappa+\kappa^2}}\right)\,.$$

Regarding τ_1, τ_2 as functions of κ, we have $\tau_2(\kappa) = 1/3$ not only in the Haar measure case $\kappa = 0$, but also for $\kappa = 1$. For the Fisher-von Mises distribution, we obtain from [4, p. 425] that

$$f^{\mathrm{FvM}}_{X}(x) = \frac{2\exp\{-2\kappa\}}{\pi\{\mathcal{I}_0(2\kappa) - \mathcal{I}_1(2\kappa)\}}\sqrt{\frac{1-x}{x}}\,\exp(4\kappa x)\,,$$

where $\mathcal{I}_0, \mathcal{I}_1$ are modified Bessel functions of the first kind, see Appendix 1 in [5], and from this we may again obtain the corresponding values of ρ_1, ρ_2 and then of τ_1, τ_2 from (9). Figure 1 illustrates the behaviour of $\tau_2(\kappa) - 1/3$, for either the Cayley-LMR or the Fisher-von Mises density, for $\kappa \in [0, 1]$. The instance $\kappa = 1$ for the Cayley-LMR distribution is of special interest and motivates the following definition.

Definition 1. Let $\mathcal{P}$ be a family of rotation distributions as described in Section 2. We say that $\mathcal{P}$ has the *fake uniformity* property if there exists $\kappa > 0$ such that the Gram matrix in the context of Theorem 1 has the same distribution as for μ.

The fake uniformity property sets the Cayley-LMR distribution distinctly apart from the Fisher-von Mises distribution in a novel aspect, besides the matter of the singular behaviour of the former at $\Theta = \pi$ already noted in [4, p. 425]. It considerably complicates, if not frustrates, any attempt at the development of a statistical test for the rotation distribution to be Haar measure, if only the random projections are known. For it to be present, a sufficient condition is that the derivative of $\tau_2(\kappa)$ at $\kappa = 0+$ is negative. By (9), the latter condition is satisfied if and only if

$$\frac{4}{3}\,\rho_2(0+) < \rho_1(0+)\,.$$

Note that this condition would not change if κ was instead reparametrised by a monotone function $\tilde{\kappa} = \varphi(\kappa)$ with $\varphi(0) = 0$.

Finally, we note that for both distributions $\tau^2(\kappa) \to 1$, and hence $D_\kappa \to \mathrm{diag}(0, 0, 1)$ and

$$\begin{aligned}\int_{SO(3)} \mathrm{Gram}(HAV) f_{\boldsymbol{P}}(A)\mu(\mathrm{d}A) &\to V^T \cdot \left[I - \mathrm{E}\left\{(Me^{(3)})(Me^{(3)})^T\right\}\right] \cdot V \\ &= \mathrm{Gram}\left\{\left(I - (Me^{(3)})(Me^{(3)})^T\right) V\right\}\end{aligned}$$

as $\kappa \to \infty$. This is intuitive: as the concentration parameter diverges to infinity, for Gram inversion, only the projection to the space orthogonal to the last column of M is relevant.



Figure 1: Values of $\tau_2(\kappa) - 1/3$ as a function of κ for the Fisher-von Mises and the Cayley-LMR distribution.

5 Bayes classifier for shift-symmetric rotations

We continue to use the notation and framework from Section 4. Consider now two shift-symmetric random rotations $I \neq \boldsymbol{P}_i \in \mathcal{I}_s^{M_i}$, $i = 1, 2$, with $M_1 \neq M_2$. There exist $Q \in O(3)$ and $\alpha \in (0, \pi)$ such that

$$I - M_1 M_2^T = Q^T A(\alpha) Q \qquad \text{with} \quad A(\alpha) = (a_{ij}(\alpha)) = \begin{pmatrix} 1-\cos\alpha & \sin\alpha & 0 \\ -\sin\alpha & 1-\cos\alpha & 0 \\ 0 & 0 & 0 \end{pmatrix}. \tag{11}$$

The angle α in (11) is the Riemannian distance between the rotations $\boldsymbol{P}_i$ when coupled as in our definitions; see [6]. Our main result in this section is as follows. For simplicity we assume equal prior probabilities $\pi_1 = \pi_2 = 0.5$ for selecting an object distributed as $\boldsymbol{P}_1$ or $\boldsymbol{P}_2$.

Theorem 2. With the foregoing notation and conditions, let T denote the event of correctly classifying a Class 1 [respectively, arbitrary] observation to Class 1 [respectively, its true class]. Then

$$\begin{aligned} \mathrm{P}(T) = {} & \mathrm{P}\left(\Theta < \frac{\alpha}{2}\right) + \frac{1}{2}\mathrm{P}\left(\frac{\alpha}{2} \leq \Theta \leq \pi - \frac{\alpha}{2}\right) \\ & + \frac{\tan\left(\frac{\alpha}{4}\right)}{2}\mathrm{E}\left\{\cot\left(\frac{\Theta}{2}\right);\, \frac{\alpha}{2} \leq \Theta \leq \pi - \frac{\alpha}{2}\right\} \\ & + \frac{1}{\sin\left(\frac{\alpha}{2}\right)}\mathrm{E}\left\{\cot\left(\frac{\Theta}{2}\right);\, \pi - \frac{\alpha}{2} < \Theta < \pi\right\}. \end{aligned}$$

Remark 3. The Gram matrix is generally unsuitable for classification. To appreciate why, take M_i such that $M_1 e^{(3)} = M_2 e^{(3)}$. The two shapes of the projections do not differ and hence classification is impossible.

Proof of Theorem 2 is at the end of this section. We first discuss classification accuracy for small α. Define

$$h(x) := \sqrt{\frac{x}{1-x}}\, f_X(x)\,, \qquad 0 \le x \le 1\,.$$

Note that for the Cayley-LMR distribution with $M = I$, we have $h(x) = x^\kappa/\mathrm{B}(\kappa + 1/2, 3/2)$. Regarding the left-hand side in Theorem 2 as a function $\psi_{\boldsymbol{R}}(\alpha)$ and writing $w_\alpha = \cos(\alpha/2)$, we have

$$\begin{aligned}\psi_{\boldsymbol{R}}(\alpha) = &\int_{\frac{1+w_\alpha}{2}}^{1} \sqrt{\frac{1-x}{x}}\, h(x)\,\mathrm{d}x + \frac{1}{2}\int_{\frac{1-w_\alpha}{2}}^{\frac{1+w_\alpha}{2}} \sqrt{\frac{1-x}{x}}\, h(x)\,\mathrm{d}x \\ &+ \frac{\tan\left(\frac{\alpha}{4}\right)}{2}\int_{\frac{1-w_\alpha}{2}}^{\frac{1+w_\alpha}{2}} h(x)\,\mathrm{d}x + \frac{1}{\sin\left(\frac{\alpha}{2}\right)}\int_0^{\frac{1-w_\alpha}{2}} h(x)\,\mathrm{d}x\,.\end{aligned}$$

It is readily checked that, with the ± signs taken respectively,

$$\sqrt{\frac{1-\frac{1\pm w_\alpha}{2}}{1\pm\frac{1\pm w_\alpha}{2}}} = \left\{\tan\left(\frac{\alpha}{4}\right)\right\}^{\pm 1}\,,$$

from which we obtain after some manipulations the simple expression

$$\psi'_{\boldsymbol{R}}(\alpha) = \frac{w_\alpha}{4(1+w_\alpha)}\left\{\frac{1}{w_\alpha}\int_{\frac{1-w_\alpha}{2}}^{\frac{1+w_\alpha}{2}} h(x)\,\mathrm{d}x - \frac{2}{1-w_\alpha}\int_0^{\frac{1-w_\alpha}{2}} h(x)\,\mathrm{d}x\right\}\,, \tag{12}$$

a weighted difference between two integral means. This expression is analogous to the well-known one for the case of two multivariate Normal distributions with equal covariance matrix given in [10, p. 22], when $\alpha = 0$. One may also regard it as a measure for deviation from uniformity, but unlike the one presented in the previous section it is a non-local one and hence probably less useful in practice.

Proof of Theorem 2. We only need to prove the assertion in the case where T is the event of correct classification into Class 1. We have

$$\mathrm{P}(\text{Observation correctly assigned to Class 1}) = \mathrm{P}\big(\mathrm{tr}\big\{\boldsymbol{P}M_1^T(I - M_1M_2^T)\big\} > 0\big)\,.$$

Moreover, with the notation at (11),

$$\mathrm{tr}\big\{\boldsymbol{P}M_1^T(I - M_1M_2^T)\big\} = \mathrm{tr}\big\{\boldsymbol{P}M_1^TQ^TA(\alpha)Q\big\} = \mathrm{tr}\big\{Q\boldsymbol{P}M_1^TQ^TA(\alpha)\big\} = \mathrm{tr}\big\{\boldsymbol{P}M_1^TA(\alpha)\big\}\,.$$

Using the representation at (2), we have

$$\begin{aligned}
&\boldsymbol{P}M_1^T = (b_{ij}) = (1-\cos\Theta)UU^T + \cos(\Theta)I + \sin(\Theta)\,\mathcal{S}\big(U\big)\,,\\
&\qquad b_{11} = (1-\cos\Theta)U_1^2 + \cos\Theta\,, \qquad\qquad b_{12} = (1-\cos\Theta)U_1U_2 - U_3\sin\Theta\,,\\
&\qquad b_{21} = (1-\cos\Theta)U_1U_2 + U_3\sin\Theta\,, \qquad\qquad b_{22} = (1-\cos\Theta)U_2^2 + \cos\Theta\,.
\end{aligned}$$

We therefore obtain

$$\begin{aligned}
\operatorname{tr}(\boldsymbol{P}M_1^T A) &= \sum_{i,j=1}^{2} b_{ij}a_{ji}(\alpha) = (1-\cos\alpha)(b_{11}+b_{22}) + (\sin\alpha)(b_{21}-b_{12})\\
&= (1-\cos\alpha)(1-\cos\Theta)(1-U_3^2) + 2(1-\cos\alpha)\cos\Theta + 2(\sin\alpha)U_3\sin\Theta\\
&= aU_3^2 + bU_3 + c\,,\\
a = -2(1-\cos\alpha)(1-X) < 0\,,&\qquad b = 4(\sin\alpha)\sqrt{X(1-X)}\,,\qquad c = 2(1-\cos\alpha)X\,.
\end{aligned} \tag{13}$$

The zeros of this quadratic in U_3 are $U_3^{(1)} \le U_3^{(2)}$ with $U_3^{(1,2)} = (2a)^{-1}\big(b \pm \sqrt{b^2-4ac}\big)$, where

$$\begin{aligned}
-\frac{b}{2a} &= \frac{\sin\alpha}{1-\cos\alpha}\sqrt{\frac{X}{1-X}} = \cot\left(\frac{\alpha}{2}\right)\cot\left(\frac{\Theta}{2}\right)\,,\\
\frac{\sqrt{b^2-4ac}}{2a} &= -\frac{\sqrt{16(\sin^2\alpha)X(1-X) + 16(1-\cos\alpha)^2X(1-X)}}{-4(1-\cos\alpha)(1-X)}\\
&= \frac{1}{1-\cos\alpha}\sqrt{\sin^2\alpha + (1-\cos\alpha)^2}\cot\left(\frac{\Theta}{2}\right) = \csc\left(\frac{\alpha}{2}\right)\cot\left(\frac{\Theta}{2}\right)\,,
\end{aligned}$$

hence (noting that $\cot\beta - \csc\beta = -\tan(\beta/2)$ and $\cot\beta + \csc\beta = \cot(\beta/2)$)

$$U_3^{(1)} = -\tan\left(\frac{\alpha}{4}\right)\cot\left(\frac{\Theta}{2}\right)\,,\qquad U_3^{(2)} = \cot\left(\frac{\alpha}{4}\right)\cot\left(\frac{\Theta}{2}\right)\,.$$

It is easily seen that $U_3^{(1)} \le -1$ if and only if $\Theta \le \alpha/2$, and $U_3^{(2)} \ge 1$ if and only if $\Theta \le \pi - \alpha/2$. Thus the integration with respect to the distribution F_{U_3} of U_3 is over the domains

$$\int_{-1}^{1} \Longleftrightarrow \Theta < \frac{\alpha}{2}\,,\qquad \int_{U_3^{(1)}}^{1} \Longleftrightarrow \frac{\alpha}{2} \le \Theta \le \pi - \frac{\alpha}{2}\,,\qquad \int_{U_3^{(1)}}^{U_3^{(2)}} \Longleftrightarrow \Theta > \pi - \frac{\alpha}{2}\,. \tag{14}$$

The density of $\arcsin U_3$ is $\{\gamma \mapsto (\cos\gamma)/2\}$ for $\gamma \in [-\pi/2, \pi/2]$. Changing variables and integrating over F_{U_3} gives for the three expressions in (14), respectively, $\mathrm{P}(\Theta \leq \alpha/2)$,

$$\int_{\alpha/2}^{\pi-\alpha/2} F_\Theta(\mathrm{d}\theta) \int_{U_3^{(1)}}^{1} F_{U_3}(\mathrm{d}u) = \frac{1}{2}\int_{\alpha/2}^{\pi-\alpha/2} \left\{1+\tan\left(\frac{\alpha}{4}\right)\cot\left(\frac{\Theta}{2}\right)\right\} F_\Theta(\mathrm{d}\theta)$$
$$= \frac{1}{2}\,\mathrm{P}\left(\frac{\alpha}{2} \leq \Theta \leq \pi - \frac{\alpha}{2}\right) + \frac{\tan\left(\frac{\alpha}{4}\right)}{2}\,\mathrm{E}\left\{\cot\left(\frac{\Theta}{2}\right);\, \frac{\alpha}{2} \leq \Theta \leq \pi - \frac{\alpha}{2}\right\},$$
$$\int_{\pi-\alpha/2}^{\pi} F_\Theta(\mathrm{d}\theta) \int_{U_3^{(1)}}^{U_3^{(2)}} F_{U_3}(\mathrm{d}u) = \frac{1}{\sin\left(\frac{\alpha}{2}\right)}\,\mathrm{E}\left\{\cot\left(\frac{\Theta}{2}\right);\, \pi - \frac{\alpha}{2} < \Theta < \pi\right\},$$

where we used the fact that

$$U_3^{(2)} - U_3^{(1)} = \left\{\tan\left(\frac{\alpha}{4}\right) + \cot\left(\frac{\alpha}{4}\right)\right\}\cot\left(\frac{\Theta}{2}\right) = \frac{2}{\sin\left(\frac{\alpha}{2}\right)}\cot\left(\frac{\Theta}{2}\right).$$

This concludes the proof of Theorem 2. □

References

[1] J. Faraut. Analysis on Lie groups: an introduction. Cambridge University Press, Cambridge, 2008.

[2] G. Jennings. Modern geometry with applications. Universitext. Springer-Verlag, New York, 1994.

[3] H. Le and D. Barden. On the induced distribution of the shape of a projection of a randomly rotated configuration. Adv. Appl. Prob., 42:331–346, 2010.

[4] C. León, J.-C. Massé, and L.-P. Rivest. A statistical model for random rotations. J. Multivariate Anal., 97:412–430, 2006.

[5] K. Mardia and P.E. Jupp. Directional statistics. Wiley Series in Probability and Statistics. John Wiley & Sons Ltd., Chichester, 2000.

[6] M. Moakher. Means and averaging in the group of rotations. SIAM J. Matrix Anal. Appl., 24:1–16, 2002.

[7] V. Panaretos. On random tomography with unobservable projection angles. Ann. Statist., 37:3272–3306, 2009.

[8] V. Panaretos and K. Konis. Sparse approximations of protein structure from noisy random projections. Ann. Appl. Statist., 5:2572–2602, 2011.

[9] C. Rau and S.N. Chiu. Grain rotations and distortions in the asymptotic variance of vacancy of the Boolean model. *J. Math. Anal. Appl.*, 384:647–657, 2011.

[10] B. Ripley. *Pattern recognition and neural networks*. Cambridge University Press, Cambridge, 2007. Reprint of the 1996 original.

[11] L.-P. Rivest and T. Chang. Regression and correlation for 3 × 3 rotation matrices. *Canad. J. Statist.*, 34:187–202, 2006.

[12] W. Schindler. *Measures with symmetry properties*, volume 1808 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2003.

[13] R. Schneider and W. Weil. *Stochastic and integral geometry*. Springer, Berlin, 2008.